



**Immunological profile of diabetic foot ulcers: update review**

Marwa Mohammed Ali Jassim<sup>1</sup>, Lana Nazar Abdul-Razzaq<sup>2</sup>, Murtada H. Hussein<sup>3\*</sup>

**Review Article**

**Abstract**

The most frequent consequence of diabetes mellitus, diabetic foot ulcers frequently do not heal and necessitate lower limb amputation. According to estimates, DFU accounts for 50% to 70% of all lower limb amputations, and 50% of patients with DFU have peripheral artery disease (PAD), which is occasionally brought on by atherosclerosis. Deformity of foot, peripheral artery disease (PAD), loss protective sensation (LOPS), and a history of foot ulcers. Global prevalence for DFUs varies by region. One of the frequent complications of diabetes in poor nations, such as Iraq, is diabetic foot; approximately two-thirds of the patients with diabetes have Diabetic Foot disorder (DFD) in Iraq and this health issue had significant negative social and economic effects. The Saudi Arabia and Bahrain have the highest prevalence rates of DFU; the reported yearly incidence of diabetic foot ulcers ranges from "2.1% to 7.4%". There are three types of diabetic foot ulcers: neuropathic, neuroischaemic, and ischemic. An infected DFU often contains three to five different types of bacteria, including gram-positive aerobes, such as "Staphylococcus sp.", gram-positive anaerobes, gram negative aerobes, gram negative anaerobes, and fungi (*Candida spp.*). Numerous studies have demonstrated that effective management of DFU can significantly decrease, postpone, or even completely avoid consequences like infection, gangrene, amputation, and death.

**Keywords:** DFU, DFD, DM, aerobic bacteria, anaerobic bacteria

\* Correspondence author: [dr-murtadahafedh@sci.utq.edu.iq](mailto:dr-murtadahafedh@sci.utq.edu.iq)

<sup>1</sup> College of Dentistry, Al-Muthanna University

<sup>2</sup> Ministry of Education, Educational Rusafa Directorate, Baghdad

<sup>3</sup> College of Science, Thi-Qar University, Thi-Qar

Received 19 October 2022, Revised 02 January 2023, Available online 11 January 2023.

Copyright © 2023 MD. This is article distributed under the terms of the Creative Commons Attribution License

<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited

## Review Article

"Diabetes mellitus" is a collection of metabolic illnesses marked by an abnormal rise in blood glucose due to impaired insulin production, insulin action, or both [1]. The most frequent complication of diabetes mellitus, diabetic foot ulcers sometimes fail to heal and result in lower limb amputation. Indeed, if the right care is not given, DFU can result in infection, gangrene, amputation, and even death [2].

DFU is thought to be the cause of 50% to 70% of all lower limb amputations [3]. It is crucial to highlight that not all diabetic individuals are at risk of developing diabetes foot ulcers (DFU), as there are several factors that contribute to them. It is estimated that approximately 26 million people globally experience diabetes foot ulcers every year [4].

Peripheral artery disease (PAD), which is a risk factor for lower limb amputations and is occasionally brought on by atherosclerosis, affects 50% of people with diabetes foot ulcers. The injury-triggered complicated biological systems that make up the wound-healing process include four essential steps: Hemostasis, inflammation, proliferation, and tissue remodeling were four factors that affected the healing of diabetic foot ulcers, along with the ulcer's location, the duration of the patient's diabetes, the age of the ulcer, the existence of heart failure, and peripheral arterial disease [5].

### Epidemiology of Diabetic Foot

Global prevalence of DFUs varies by region. In Iraq, the prevalence of diabetes mellitus in our country was 9.3% in 2015 according to statistics published on International Diabetes Federation [6]. One of the frequent complications of diabetes in impoverished nations, such as our "own country". Approximately two-thirds of the patients with diabetes have Diabetic Foot disorder (DFD) in Iraq and This health issue had significant negative social and economic effects. In study by [7] in Babylon, Iraq, illustrated that incidence rate was increasing and more prominent among urban-dwelling, low-educated women. As well as, in study by [8] in Basrah, Iraq showed that DFD were strongly associated with long duration of diabetes and the female gender. According to a study by [9], the prevalence rates of DFU were lowest in Iraq and highest in Saudi Arabia and Bahrain.

The lifetime risk of getting a diabetic foot ulcer has been estimated to be as high as 25%, and the reported annual incidence of diabetic foot ulceration ranges from 2.1% to 7.4% (10). It has been estimated that up to 15% of persons with T2DM have DFUs [11]. Globally, the incidence is estimated to be 6.3% (95%) with a reported 13.0% prevalence in North America. 5.5% is cited as the prevalence in Asia; in Europe it is 5.1 %, in Africa it is 7.2 % and in the Oceania region it is 3.0 %. Additionally, individuals with DM for longer than ten years are more prone

to acquire a DFU [12]. Amputation is the diabetic foot ulcer's final outcome in 15% to 27% of cases if it is not treated promptly and effectively [13].

Additionally, amputation typically has negative social, psychological, and financial effects in addition to high morbidity and mortality [14].

### Classification of diabetes foot ulcers:

There are three types of diabetic foot ulcers: neuropathic, neuroischemic, and ischemic [15]. In order to describe the severity of an ulcer, doctors also use the Wagner Grades. The Wagner Grades are designed to help professionals better monitor and care for diabetic foot ulcers. This grading scale uses numbers from 0 to 5, as shown in the table, to categorize diabetic foot ulcers [1-2].

**Table 1.**

Classification of diabetic foot ulcers suggested via Wagner (Wagner *et al.*, 1983) [16]

Grade-0	No ulcer in a high-risk foot.
Grade - I	Superficial ulcer involving the full skin thickness but not underlying tissues.
Grade - II	Full-thickness ulcers, penetrating through fat to tendon, or joint capsule without causing a deep abscess or osteomyelitis.
Grade - III	Deep ulcer with abscess formation, often with osteomyelitis.
Grade - IV	Localized gangrene.
Grade - V	Extensive gangrene that involves the entire foot

### Microbial Etiology of Diabetic Foot Ulcer

Aerobes and anaerobic microorganism usually cause diabetic foot wounds to get infected, and this infection subsequently results in the production of microthrombi, which causes ischemia, necrosis, and progressive gangrene before being amputated [17]. Patients with diabetic foot have reduced microvascular circulation, which restricts phagocyte availability and encourages the development of infection.

Several of studies illustrated at present a wide variety of microorganisms discovered from infections in diabetic feet, demonstrating the long-lasting, exposed, and anatomical position of these infections. Both mono- and polymicrobial DFIs are possible, with polymicrobial DFIs being more prevalent in chronic illnesses that have already received antibiotic treatment. Human skin is home to a wide variety of pathogenic and non-pathogenic microorganisms.

An infected DFU often contains three to five different types of bacteria, including the following: gram-positive anaerobes (Enterococcus species, Propionibacterium species, Streptococcus species, Peptostreptococcus species, Peptococcus species), gram-negative aerobes (Pseudomonas aeruginosa, Acinetobacter species), gram-negative anaerobes (Proteus mirabilis, Escherichia coli, Bacteroides species), and (Candida spp.). In low-income countries, there is a higher prevalence of gram-negative pathogens, the most common of which is Pseudomonas aeruginos. Streptococcus and gram-positive cocci, in particular Staphylococci, are frequently isolated [18,19, 20].

Some investigations have revealed that the presence of anaerobic microbes is related to deeper DFIs. Diabetes patients are more prone to foot infections due to neuropathy, vascular dysfunction, and lowered neutrophil activity. A foot infection develops mostly as a result of peripheral neuropathy, which affects between 30 and 50 percent of diabetic individuals [21]. In addition, bacteria typically create biofilms that thwart immune clearance and encourage the development of antimicrobial resistance; in one research, 78.2% of chronic wounds exhibited biofilm creation. To guide the development of innovative therapies and inform antimicrobial therapy, a deeper comprehension of the microbiology of DFIs is crucial [22, 23].

### **Human and animal rights**

Non

### **Consent for publication**

Non

### **Conflict of interest**

The authors declare no conflict of interest, financial or otherwise.

### **References**

1. Seidu S, Cos X, Brunton S, et al. 2022 update to the position statement by Primary Care Diabetes Europe: a disease state approach to the pharmacological management of type 2 diabetes in primary care. Primary Care Diabetes. 2022. <https://doi.org/10.1016/j.pcd.2022.02.002>
2. Buse JB, Wexler DJ, Tsapas A, et al. 2019 update to: management of hyperglycemia in type 2 diabetes, 2018. A consensus report by the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). Diabetes care. 2020; 43(2):487-493. <https://doi.org/10.2337/dci19-0066>
3. Liu R, Li L, Shao C, et al. The Impact of Diabetes on Vascular Disease: Progress from the Perspective of Epidemics and Treatments. Journal of Diabetes Research, 2022. <https://doi.org/10.1155/2022/1531289>

4. Adams JD, Egan AM, Laurenti MC, et al. Insulin secretion and action and the response of endogenous glucose production to a lack of glucagon suppression in nondiabetic subjects. *American Journal of Physiology-Endocrinology and Metabolism* 2021; 321(5): E728-E736.  
<https://doi.org/10.1152/ajpendo.00284.2021>
5. Aldhaefi M, Aldardeer NF, et al. Updates in the Management of Hyperglycemia Crisis. *Frontiers in Clinical Diabetes and Healthcare*. 2022; 35. <https://doi.org/10.3389/fcdhc.2021.820728>
6. Tian X, Zuo Y, Chen S, et al. Hypertension, Arterial Stiffness, and Diabetes: a Prospective Cohort Study. *Hypertension*. 2022; 10-1161.
7. Jiang DH, O'Connor PJ, Huguet N, et al. Modernizing Diabetes Care Quality Measures: Analysis examines modernizing diabetes care quality measures. *Health Affairs*. 2022; 41(7):955-962.  
<https://doi.org/10.1377/hlthaff.2022.00233>
8. El-Kebbi IM, Bidikian NH, Hneiny L, et al. Epidemiology of type 2 diabetes in the Middle East and North Africa: Challenges and call for action. *World journal of diabetes*. 2021; 12(9):1401.  
<https://doi.org/10.4239/wjd.v12.i9.1401>
9. Hamdy O, Al Sifri S, Hassanein M, et al. The Transcultural Diabetes Nutrition Algorithm: A Middle Eastern Version. *Frontiers in Nutrition*. 2022; 9. <https://doi.org/10.3389/fnut.2022.899393>
10. Antal B, McMahon LP, Sultan SF, et al. Type 2 diabetes mellitus accelerates brain aging and cognitive decline: Complementary findings from UK Biobank and meta-analyses. *Elife*. 2022; 11:e73138.  
<https://doi.org/10.7554/eLife.73138>
11. Al-Shaheeb H, Hashim K, Mohammed AK, et al. Assessment of lipid profile with HbA1c in type 2 diabetic Iraqi patients. *Revis Bionatura* 2022; 7(3):29. <https://doi.org/10.21931/RB/2022.07.03.29>
12. Mathur P, Leburu S, Kulothungan V. Prevalence, awareness, treatment and control of diabetes in India from the countrywide National NCD Monitoring Survey (NNMS). *Frontiers in public health*. 2022: 205. <https://doi.org/10.3389/fpubh.2022.748157>
13. Mitsugashira H, Imura T, Inagaki A, et al. Development of a novel method for measuring tissue oxygen pressure to improve the hypoxic condition in subcutaneous islet transplantation. *Scientific Reports*. 2022; 12(1):1-11. <https://doi.org/10.1038/s41598-022-19189-2>
14. Colclough K, Ellard S, Hattersley A, Patel K. Syndromic monogenic diabetes genes should be tested in patients with a clinical suspicion of maturity-onset diabetes of the young. *Diabetes*. 2022; 71(3):530-537. <https://doi.org/10.2337/db21-0517>
15. Ertuğrul B, Uçkay I, Schöni M, Peter-Riesch B, Lipsky BA. Management of diabetic foot infections in the light of recent literature and new international guidelines. *Expert Review of Anti-infective Therapy*. 2020; 18(4):293-305.
16. Wagner FW, O'Neal LW. *The Diabetic Foot*, Mosby, St Louis. p.274. Brodsky JW. Outpatient diagnosis and care of the diabetic foot. *J. Instr Course Lect*. 1983; 42:121.  
<https://doi.org/10.1080/14787210.2020.1730177>
17. Parhi A, Das S, Mahapatra S, et al. The Level and Role of Interleukin-17 in Patients of Type 2 Diabetes Mellitus with and without Complications. *Journal of diabetes mellitus*. 2019; 9(04):176.
18. Bekele F, Chelkeba L, Fekadu G, Bekele K. Risk factors and outcomes of diabetic foot ulcer among diabetes mellitus patients admitted to Nekemte referral hospital, western Ethiopia: Prospective observational study. *Annals of Medicine and Surgery*. 2020; 51:17-23.

19. Djergovic D, Xu Z, Barrat FM, Xhing C. G-ODN protects diabetes mellitus complicated with cerebral infarction in mice model. *American Journal of BioMedicine* 2016;4(11):480-504.  
<https://doi.org/10.18081/2333-5106/016-11/480-504>
20. Tola A, Regassa LD, Ayele Y. Prevalence and associated factors of diabetic foot ulcers among type 2 diabetic patients attending chronic follow-up clinics at governmental hospitals of Harari Region, Eastern Ethiopia: A 5-year (2013–2017) retrospective study. *SAGE open medicine*. 2021; 9: 2050312120987385. <https://doi.org/10.1177/2050312120987385>
21. Kaminski MR, Lambert KA, Raspovic A, et al. Risk factors for foot ulceration in adults with end-stage renal disease on dialysis: a prospective observational cohort study. *BMC nephrology* 2019; 20(1):1-11. <https://doi.org/10.1186/s12882-019-1594-5>
22. Liu J, Yuan X, Liu J, et al. Risk Factors for Diabetic Peripheral Neuropathy, Peripheral Artery Disease, and Foot Deformity Among the Population with Diabetes in Beijing, China: A Multicenter, Cross-Sectional Study. *Frontiers in endocrinology*. 2020; 13. <https://doi.org/10.3389/fendo.2022.824215>
23. Pereira M, Oh JK, Kang DK, et al. Hacking Commensal Bacteria to Consolidate the Adaptive Mucosal Immune Response in the Gut–Lung Axis: Future Possibilities for SARS-CoV-2 Protection. *BioTech*. 2022; 11(1):3. <https://doi.org/10.3390/biotech11010003>