

<http://dx.doi.org/10.52113/1/1/2024-2-81>

Overview on the most common microbial causative agents of diarrhea

Wafaa Ayad Al-Nuaimy ¹, Dhay A. Azeez ¹, Shaimaa MS. Zainulabdeen ², Iman Ali Atiyah ³



Abstract

In all parts of the world and among all age groups, diarrhea is a significant cause of morbidity and mortality. However, compared to high-income countries, low-income countries have fewer resources and a less resilient infrastructure, which makes them more susceptible to an unfair share of diarrheal morbidity and mortality. The word "diarrhea" originates from the Greek word "dia rEo," referring to "flow through." When the volume of colonic fluid crosses this segment's absorptive capacity due to decreased absorption and/or increased secretion, diarrhea achievements. In infectious diarrheas, the abnormal function is brought about by microorganisms that colonize the intestinal mucosa and subvert normal gut physiology either directly or via enterotoxins.

Keywords: Morbidity and mortality, Diarrheal, Colonic fluid

* Correspondence author: wafaa.ayad@mu.edu.iq

¹ College of Science, Al-Muthanna University, Iraq

² College of Medicine, Al-Muthanna University, Iraq

³ Department of Medical Laboratories, College of Health and Medical Technology, Sawa University, Iraq.

Received 21 May 2024; revised 18 June 2024; accepted 11 July 2024, available online 01 August 2024.

Copyright © 2024 Wafaa, et al. This is article distributed under the terms of the Creative Commons Attribution License <http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited

Introduction

The passage of loose or watery stools that tend to be accompanied by an increase in the frequency of bowel movements is termed as diarrhea. It can also mean passing no fewer than three watery or loose stools in a 24-hour period; a loose stool conforms to the shape of the container [1, 2, 3]. Children under the age of two years old have higher rates of diarrheal illness in developing nations. Under-five children experience 4 million deaths and 1.3 billion episodes of diarrhoea annually, with an average of 4.9 episodes of diarrhoea / child a year [3]. Globally, diarrheal diseases are a leading cause of morbidity and mortality, particularly in developing nations where children are most affected by infectious diarrhea, malnourishment, and death. Diarrheal diseases claim the lives of about 5 million children and infants worldwide each year [4].

In children acute diarrhea is primarily caused by infectious organisms, such as viruses, bacteria, and parasites. Since parasite infections are now less common, viruses and bacteria are now more commonly to blame for severe diarrhea in kids [5, 6]. One of the main causes of diarrhea in children, particularly in those under five, is the human rotavirus. The majority of global reports concur that rotavirus is the main culprit behind children's acute diarrhea [7,8]. However, it seems that the etiology of the bacteria that cause diarrhea varies by region. For instance, a study from Spain found that the

main bacterial pathogens responsible for 22.2 % and 16.4% of pediatric episodes of acute diarrhea, respectively, were *Campylobacter* spp. and *Salmonella* spp. [9]. According to reports, the primary etiological causes of diarrhea in Ecuador are *Shigella* spp. and *Campylobacter jejuni* [10]. In Turkey, the primary causes of acute gastroenteritis in children are *Salmonella* spp. (25.6 %) and *C. jejuni* (18.3%) [11].

Etiology of Diarrhea:

Bacterial Pathogens

AIDS patients experience bacterial enteritis more frequently and to a greater extent than the general population. Over 20% of diarrhea episodes in AIDS patients are caused by bacterial pathogens. These pathogens include diarrhoeagenic *Escherichia coli* [which include enteropathogenic *E. coli* (EPEC), entero-invasive *E. coli* (EIEC), enteroaggregative *E. coli* (EAaggEC), entero-toxigenic *E. coli* (ETEC) [12,13], and attaching and effacing *E. coli* (A/EEC)], *Shigella* species, *Salmonella* species, *Vibrio cholera*, *Campylobacter jejuni*, *Yersinia enterocolitica*, *Clostridium difficile* and *Aeromonas* species [14,15].

Viral gastroenteritis is more common than bacterial gastroenteritis. Gastroenteritis is caused by bacteria in a few different ways.

Enterotoxins are produced by certain species (eg, *Vibrio cholerae*, enterotoxigenic strains of *E. coli*) that adhere to intestinal mucosa without invading. Watery diarrhea is the result of these toxins' inhibition of intestinal absorption and stimulation of adenylate cyclase, which leads to the secretion of water and electrolytes. A comparable toxin is produced by *C. difficile*.

Exotoxins that are ingested in contaminated food which are produced by some bacteria (eg, *Staphylococcus aureus*, *Clostridium perfringens*, *Bacillus cereus*). In the absence of a bacterial infection, the exotoxin can induce gastroenteritis. After consuming contaminated food, these toxins typically result in vomiting, acute nausea and diarrhea within 12 hours. The symptoms go away in three days.

Mucosal invasion happens when other bacteria, such as *Shigella*, *Salmonella*, *Campylobacter*, *C. difficile*, and some subtypes of *E. coli*, infiltrate the mucosa of the colon or small intestine and produce ulcers, bleeding, fluid that is rich in protein, and the secretion of electrolytes and water. Whether the organism produces an enterotoxin or not does not affect the outcome of the invasive process. There is evidence of this invasion and inflammation in the ensuing diarrhea, with leukocytes and RBCs visible under a microscope and occasionally with gross blood [16, 17].

Infections with bacterial causes such as *Salmonella* and *Campylobacter* are prevalent of diarrhea in the United States. Undercooked poultry is the most common way to contract both infections, though unpasteurized milk can also be a source. Cattle and poultry are common sources of *Campylobacter*; other animals, such as puppies and kittens, pets, pigs, and birds, may also harbor the bacteria and spread the infection to people. Acute, mostly self-limited gastrointestinal illness caused by a *Campylobacter* infection is characterized by diarrhea, fever, and cramping in the abdomen [18].

One of the three most important causes of inflammatory diarrhea in the world, *salmonella* primarily affects people who live in places without access to clean water and food, basic healthcare, appropriate nutrition, and good hygiene [19]. Undercooked eggs can spread *Salmonella*, as can coming into

contact with birds, amphibians, or reptiles. Although foodborne outbreaks do happen, *Shigella* species are frequent bacterial causes of diarrhea in United State and are typically spread from man to man. Shiga toxin produced by *Shigella dysenteriae* type 1 can brought on Hemolytic-uremic syndrome, which is, which is not found in the US [20, 21].

Many subtypes of the bacteria can cause gastroenteritis due to *E. coli*.

Depending on the subtype, the epidemiology and clinical manifestations differ significantly.

The most frequent cause of infectious nosocomial diarrhea is toxin-producing *Cl. difficile* infection, which accounts for 15% to 20% of cases of antibiotic-associated diarrhea. Furthermore, its occurrence has been increasing in recent years. More than ninety percent of cases of *Cl.difficile* diarrhea are acquired in the hospital, whereas fewer than 5 % are community-acquired. Other risk factors for *Cl.difficile* diarrhea are older age, , renal failure, severe underlying illness, enteral feedings and use of rectal thermometers. 3'4>11 There is a linear relationship between length of hospital stay, colonization with *Cl.difficile*, and development of *Cl.difficile* diarrhea [23, 24, 25].

Although most are rare in the US, a few other bacteria can also cause gastroenteritis. *Y. enterocolitica* causes acute diarrhea in early childhood. It is transmitted by undercooked pork, unpasteurized milk, or contaminated water [26]. When undercooked seafood is consumed, a number of *Vibrio* species, such as *V. parahaemolyticus*, can cause diarrhea. Particularly after natural disasters or in camp over refugees, Because it can occasionally cause severe watery diarrhea in places where people lack access to clean drinking water and sanitary ways to dispose of human waste, *V. cholerae* is a concern. There are hundreds of thousands of cholera cases reported each year, with an approximate 1-2% overall case mortality rate. Cholera toxin (CT), which is made up of five binding (B) subunits and an enzymatic (A) subunit, is the main virulence factor of *V. cholerae*. When CT is internalized and its A subunit is released into the cytosol, it stimulates intracellular cAMP production, allowing cAMP-mediated intestinal fluid secretion.

This occurs after CT's B subunit binds to GM1 ganglioside receptors found in the apical membrane of intestinal epithelial cells (IEC) [27]. Rarely can *listeria* cause foodborne gastroenteritis; instead, it typically causes meningitis or bloodstream infection in older adults, pregnant women, and newborns (see Neonatal Listeriosis). *Aeromonas* can be contracted by swimming in or drinking contaminated fresh or brackish water [28].

Viral Pathogens

Viruses are a major global cause of gastrointestinal issues, particularly affecting children under the age of five in both developed and developing nations. Adenovirus is another microbe that can cause acute diarrhea, particularly in young children and infants, along with rotavirus. Enteric adenovirus or astrovirus are the main causes of the majority of other viral gastroenteritis infections [29].

Viral gastroenteritis is frequently caused by astroviruses in both geographical and developmental contexts [30]. Seasonal outbreaks of infection are linked to hospitals, childcare centers, and hotels in temperate climates. in tropical environments with inadequate infrastructure for water and sanitation, In low- and middle-income countries (LMIC), diarrhea in children is a major cause of mortality and wasted human potential. Astroviruses are part of the problem. They are thought to be responsible for 2% to 9% of cases of acute, nonbacterial childhood diarrhea, and new strains have

been connected to complications for the central nervous system in children, such as encephalitis and meningitis. Common symptoms of astrovirus gastroenteritis include sudden, self-limiting, acute watery diarrhea; fever, anorexia, and vomiting are not always present. The mode of transmission is the fecal-oral route. For incubation, three to four days need to be spent [31, 32, 33].

Adenoviruses are the fourth most frequent cause of viral gastroenteritis in young people. All year long, infections happen, though they do tend to spike in the summer.

Epidemiological investigations conducted across multiple developed nations revealed that enteric adenovirus was only the second most frequent reason why infants and young children get viral gastroenteritis children, and its positive infection rate in isolated instances and acute gastroenteritis outbreaks varied from 1.1% to 12.0%. Youngsters under the age of two are the most vulnerable.. Both respiratory droplets and the fecal-oral pathway are modes of transmission, 3-5 days are needed for incubation [34].

Rotavirus gastroenteritis is a serious global public health issue. The feco-oral route is how rotavirus is spread, and winter is when rotavirus gastroenteritis is most common. Its clinical course is generally thought to be more severe than that of other viral gastroenteritis. A mild fever, vomiting, and loose stools are the first symptoms; there is no visible blood or mucus in the stool. Diarrhea usually lasts 4–5 days, while vomiting can last 2–3 days [35, 36].

Enterocytes in the small intestine's villous epithelium are infected by viruses. Fluid and electrolytes transude into the intestinal lumen as a result; occasionally, unabsorbed carbohydrates from malabsorption in the impacted colon subsequently exacerbate by resulting in osmotic diarrhea symptoms. Watery diarrhea occurs. Diarrhea (dysentery) characterized by inflammation and fecal red blood cells (RBCs), white blood cells (WBCs), or gross blood. Other viruses, such as the enterovirus and cytomegalovirus, can induce gastroenteritis in immunocompromised patients [37].

Protozoan Pathogens

In humans, protozoan parasites are major contributors to diarrhea and other intestinal disorders [38]. In developing nations, *Giardia lamblia* and *Cryptosporidium parvum* are two significant protozoan causes of diarrhea. Youngsters under two years old may be the most common, and they may also be the most vulnerable. The associated diseases have a wide range of clinical manifestations, from self-limited diarrhea in healthy individuals and asymptomatic carriers to potentially fatal prolonged diarrhea in immunocompromised or immunosuppressed individuals (such as those with AIDS), malnourished children, and the elderly) [39, 40].

Giardia and *Cryptosporidium* spread by tainted food and water, but they can also spread from person to person and, because these protozoans are zoonotic, from domestic or wild animals to humans. Despite being acknowledged as important foodborne pathogens, foodborne parasites are still given less attention than bacterial and viral pathogens [41].

Some intestinal parasites stick to the intestinal mucosa and cause nausea, vomiting, diarrhea, and general malaise. One such parasite is *Giardia intestinalis* (also known as *G. lamblia*). Globally and in every US region, giardiasis is a problem. Chronic infections can lead to malabsorption syndrome, which is sometimes mistaken for irritable bowel syndrome. Typically, people contract it through person-to-person contact (common in daycare facilities) or by ingesting cysts in tainted food or water.

In addition to causing watery diarrhea, *Cryptosporidium parvum* can occasionally cause nausea, vomiting, and cramping in the abdomen. In healthy individuals, the illness resolves on its own after about two weeks. Immunocompromised patients may experience a severe and protracted illness that results in significant fluid and electrolyte loss. The most common way to get cryptosporidium is through contaminated water. Roughly three-quarters of outbreaks of recreational waterborne illness in the US are caused by this organism, which is resistant to chlorine [42, 43].

Entamoeba histolytica is a common cause of subacute bloody diarrhea in areas with low socioeconomic status and inadequate sanitation, despite being uncommon in the US. The surrounding environment, including social, economic, demographic, and hygiene-related behaviors that affect the spread and transmission of parasitic infections, are important in determining the prevalence of *E. histolytica*. A numerous studies identified that drinking water quality, ingestion of raw vegetables, place of residence and age as important risk factors [44]. Some researchers conjectured that infants should not be expected to contract amebiasis very often because the disease is often linked to contaminated food and water. Malnutrition, immunization, and young age are linked to more severe disease [45].

Other parasites that can mimic the symptoms of cryptosporidiosis include a group of organisms known as *microsporidia* and *Cystoisospora (Isospora) belli*. The gastrointestinal tract is home to the coccidian, unicellular protozoan parasite *I. belli*. Although *I. belli* infections are found all over the world, they are comparatively uncommon and typically arise in tropical and subtropical regions. It typically causes diarrhea that isn't bloody in tropical and subtropical regions. It is present in newly arrived immigrants, tourists coming back from endemic areas, and AIDS patients in developed nations. The most common way for *I. belli* to spread is through contaminated food or water [46, 47].

Conflict of Interest

No conflicts of interest were declared by the authors.

Financial Disclosure

The authors declared that this study has received no financial support.

Ethics Statement

Approved by local committee.

Authors' contributions

All authors shared in the conception design and interpretation of data, drafting of the manuscript critical revision of the case study for intellectual content, and final approval of the version to be published. All authors read and approved the final manuscript.

Open access

This is an open-access article distributed by the Creative Commons Attribution Non-Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial.

<http://creativecommons.org/licenses/by-nc/4.0/>.

References

1. Odimayo MS, Fowotade TA, Adegboro B. Opinion of Care givers on the possible cause of Diarrhea among children in Ilorin, Nigeria. *Pioneer Medical Journal*. 2011;1(2):10-13.
2. Angela Ine Frank-Briggs. Introduction and Classification of Childhood Diarrhoea, Current Concepts in Colonic Disorders, Dr. Godfrey Lule (Ed.), InTech, World Health Organization. The treatment of diarrhoea. A manual for physicians and other senior health workers. 2005.
3. Silvia resta-lenert. Diarrhea Infectious, *Encyclopedia of Gastroenterology*, Elsevier (USA). 2004; 576-584.
4. Kotloff KL, Nataro JP, Blackwelder WC, Nasrin D, et al. Burden and aetiology of diarrhoeal disease in infants and young children in developing countries (the Global Enteric Multicenter Study, GEMS): a prospective, case-control study. *Lancet*. 2013;382(9888):209–22.
5. Lanata CF, Fischer-Walker CL, Olascoaga AC, et al. Global causes of diarrheal disease mortality in children <5 years of age: a systematic review. *PLoS One*. 2013;8(9):e72788.
6. Walker CLF, Rudan I, Liu L, Nair H, Theodoratou E, Bhutta ZA, O'Brien KL, Campbell H, Black RE. Global burden of childhood pneumonia and diarrhoea. *Lancet*. 2013;381(9875):1405–16.
7. Yu J, Jing H, Lai S, Xu W, et al. Etiology of diarrhea among children under the age five in China: Results from a five-year surveillance. *J infect*. 2015;71(1):19–27.
8. Sanchez-Capilla AD, Sorlozano-Puerto A, Rodriguez-Granger J, Martinez-Brocal A, Navarro-Mari JM, Gutierrez-Fernandez J. Infectious etiology of diarrheas studied in a third-level hospital during a five-year period. *Rev Esp Enferm Dig: organo ofic Soc Esp Patol Dig*. 2015;107(2):89–97.
9. G, Trueba G, Atherton R, Calvopina M, Cevallos W, Andrade T, Eguiguren M, Eisenberg JN. Identifying etiological agents causing diarrhea in low income Ecuadorian communities. *AmJTrop Med Hyg*. 2014;91(3):563–9.
10. Bicer S, Col D, Erdag GC, Giray T, Gurol Y, Yilmaz G, Vitrinel A, Ozelgun B. A retrospective analysis of acute gastroenteritis agents in children admitted to a university hospital pediatric emergency unit. *Jundishapur j microbiol*. 2014;7(4):e9148.

11. Hien BT SF, Cam PD, Serichantalergs O, Huong TT, Thu TM, Dalsgaard A. Diarrheagenic *Escherichia coli* and *Shigella* strains isolated from children in a hospital casecontrol study in Hanoi, Vietnam. *J Clin Microbiol* 2008;46(3):996-1004.
12. El-Sheikh SM e-AS. Prevalence of viral,bacterial and parasitic enteropathogens among young children with acute diarrhoea in Jeddah,Saudi Arabia. *J Health Popul Nutr*2001;19(1):25-10.
13. Nitiema LW NJ, Ouermi D, Dianou D, TraoreAS, Svensson L, Simpore J. Burden of rotavirusand other enteropathogens among children withdiarrhea in Burkina Faso. *Int J Infect Dis* 2011;15(9):e646-52.
14. Laham NA EM, Al-Haddad R, Ridwan F.Prevalence of enteric pathogen-associated community gastroenteritis among kindergarten children in Gaza. *J Biomed Res* 2015;29(1):61-8.
15. Centers for Disease Control and Prevention (CDC): *Shigella—Shigellosis: Questions & Answers*. Accessed May 12, 2023.
16. Johnson S, Lavergne V, Skinner AM, et al. Clinical practice guideline by the Infectious Diseases Society of America (IDSA) and Society for Healthcare Epidemiology of America (SHEA): 2021 focused update guidelines on management of *Clostridioides difficile* infection in adults. *Clin Infect Dis* ciab549, 2021.
17. Heymann, D.L. *Control of Communicable Diseases Manual*, 20th ed.; American Public Health Association, Alpha Press: Washington, DC, USA, 2014.
18. Pfeiffer ML. Dupont, HL, Ochoa, T. J. The patient presenting with acute dysentery - a systematic review. *J Infect*, v. 64, p. 374-86, 2012.
19. Sakran W, Hexner-Erlichman Z, Spiegel R, et al. *Campylobacter* gastroenteritis in children in north-eastern Israel comparison with other common pathogens. *Sci. Rep.* 2020, 10, 1–5.
20. Shim JO, Chang JY, Kim A, Shin S. Different Age Distribution between *Campylobacteriosis* and Nontyphoidal *Salmonellosis* in Hospitalized Korean Children with Acute Inflammatory Diarrhea. *J. Kor. Med. Sci.* 2017;21:202–1206
21. Kaneko T, Matsuda R, Taguri ME, et al. *Clostridium difficile* infection in patients with ulcerative colitis: Investigations of risk factors and efficacy of antibiotics for steroid refractory patients. *Clin Res Hepatol Gastroenterol* 2011; 5:315-20. 6.
22. Berg AM, Kelly CP, Farraye FA. *Clostridium difficile* infection in the inflammatory bowel disease patient. *Inflamm Bowel Dis* 2013;19:194-204. 7.
23. Mylonaki M, Langmead L, Pantas A, Johnson F, Rampton DS. Enteric infection in relapse of inflammatory bowel disease: importance of microbiological examination of stool. *Eur J Gastroenterol Hepatol* 2004;16:775-8. 25.
24. Deshpande A, Pant C, Pasupuleti V, Rolston DD, Jain A, Deshpande N. Association between proton pump inhibitor therapy and *Clostridium difficile* infection in a meta-analysis. *Clin Gastroenterol Hepatol* 2012;10:225-33.
25. Rosner BM, Stark K, Hohle M, Werber D. Risk factors for sporadic *Yersinia enterocolitica* infections, Germany 2009–2010. *Epidemiol Infect* 2012;140:1738–47.

26. Muanprasat C, Chatsudhipong V. Cholera: pathophysiology and emerging therapeutic targets. *Future Med Chem* 2013; 5:781-98
27. Chu YW, Wong CH, Tsang GKL, Kwok MSW, Wong RKO, Lo JYC, Kam KM: Lack of association between presentation of diarrheal symptoms and faecal isolation of *Aeromonas* spp. amongst outpatients in Hong Kong. *J Med Microbiol* 2006; 55: 349–351.
28. Flomenberg, Morven Mary M. Diagnosis and treatment of adenovirus infection in children. *Pediatr J* 2005; 24(6):
29. Bosch A, Pintó RM, Guix S. Human astroviruses. *Clin Microbiol Rev.* 2014;27(4):1048–1074
30. Krishnan T. Novel human astroviruses: challenges for developing countries. *Virusdisease.* 2014;25(2):208–214
31. Naccache SN, Peggs KS, Mattes FM, et al. Diagnosis of neuroinvasive astrovirus infection in an immunocompromised adult with encephalitis by unbiased next-generation sequencing. *Clin Infect Dis.* 2015;60(6):919–923.
32. Herrmann JE, Taylor DN, Echeverria P, Blacklow NR. Astroviruses as a cause of gastroenteritis in children. *N Engl J Med.* 1991;324(25):1757–1760
33. Adhikary AK, Numaga J, Kaburaki T, et al. Genetic characterisation of adenovirus type 8 isolated in Hiroshima city over a 15 year period. *J. Clin. Pathol.* 2003; 56:120–125.
34. Akan, H., Izbirak, G., Gurol, Y., Sarikaya, S., et al. (2009) Rotavirus and Adenovirus Frequency among Patients with Acute Gastroenteritis and Relationship to Clinical Parameters: A Retrospective Study in Turkey. *Asia Pacifica Family Medicine*, 8, 8.
35. Tagbo, B.N., Mwenda, J.M., Armah, G., Obidike, E.O., Okafor, U.H., Oguonu, T., et al. (2014) Epidemiology of Rotavirus Diarrhea among Children Younger than 5 Years in Enugu, South East Nigeria. *The Pediatric Infectious Disease Journal*, 33, S19-S22
36. Biswajyoti, B., Aniruddha, J., Chandrakanta B., Das, M., et al. (2016) Prevalence of Enteric Adenovirus among Non-Rotavirus Diarrhoea in Assam, Northeast India. *International Journal of Medical Research Professionals*, 2, 124-130
37. Custodio, H. Protozoan parasites. *Pediatr. Rev.* 2016, 37, 59–71.
38. Mmbaga, B.T.; Houpt, E.R. Cryptosporidium and Giardia Infections in Children: A Review. *Pediatr. Clin. N. Am.* 2017, 64, 837–850.
39. Desai NT, Sarkar R, Kang G. Cryptosporidiosis: An under-recognized public health problem. *Trop. Parasitol.* 2012, 2, 91–98.
40. Cacciò SM, Thompson RA, McLauchlin J, Smith HV. Unravelling Cryptosporidium and Giardia epidemiology. *Trends Parasitol.* 2005, 21, 430–437.
41. van der Giessen J, Deksne G, Gómez-Morales MA, et al. Surveillance of foodborne parasitic diseases in Europe in a One Health approach. *Parasite Epidemiol. Control* 2021, 13, e00205.
42. O’Leary JK, Sleator RD, Lucey B. Cryptosporidium spp. diagnosis and research in the 21st century. *Food Waterborne Parasitol.* 2021, 24, e00131
43. Benetton ML, Goncalves AV, Meneghini ME, Silva EF, Carneiro M. Risk factors for infection by the *Entamoeba histolytica*/E. dispar complex: an epidemiological study conducted in

outpatient clinics in the city of Manaus, Amazon Region, Brazil. *Trans R Soc Trop Med Hyg.* 2005;99:532-40

44. Hotez, P.J., Kamath, A. 2009. Neglected tropical diseases in sub-Saharan Africa: Review of their prevalence, distribution and disease burden. *PLoS Neglected Tropical Diseases* 3:8.
45. Gruz F, Fuxman C, Errea A, et al. *Isospora belli* infection after isolated intestinal trnsplant. *Transpl Infect Dis* 2009; 12: 69-72.
46. 47. Klasse-Fischer MK, Neafie RC, Wear DJ, Meyers WM. Chapter 13. Cryptosporidiosis, isosporiasis, cyclosporiasis, and sarcocystosis. In *Topics on the Pathology of Protozoan and Invasive Arthropod Diseases (e-book)*. Fairfax, Virginia, USA. Inova Central Laboratory. 2011, p 1-19.